

THE NUCLEON “TENSOR CHARGES” AND THE SKYRME MODEL ^{*}

James M. Olness [†]

*Center for Theoretical Physics
Laboratory for Nuclear Science
and Department of Physics
Massachusetts Institute of Technology
Cambridge, MA 02139*

Submitted to: Physical Review D (Brief Reports)

CTP#2122

July 1992

^{*} This work is supported in part by funds provided by the U.S. Department of Energy (D.O.E.) under contract #DE-AC02-76ER03069.

[†] National Science Foundation Graduate Fellow

ABSTRACT

The lowest moment of the twist-two, chiral-odd parton distribution $h_1(x)$ of the nucleon can be related to the so-called “tensor charges” of the nucleon. We consider the tensor charges in the Skyrme model, and find that in the large- N_c , SU(3)-symmetric limit, the model predicts that the octet isosinglet tensor charge, g_T^8 , is of order $1/N_c$ with respect to the octet isovector tensor charge, g_T^3 . The predicted F/D ratio is then $1/3$, in the large- N_c limit. These predictions coincide with the Skyrme model predictions for the octet *axial* charges, g_A^8 and g_A^3 . (The prediction $F/D = 1/3$ for the axial charges differs from the commonly quoted prediction of $5/9$, which is based on an inconsistent treatment of the large- N_c limit.) The model also predicts that the singlet tensor charge, g_T^0 , is of order $1/N_c$ with respect to g_T^3 .

The nucleon has three parton distributions at lowest twist, that is, at twist-two. Two of these, $f_1(x)$ and $g_1(x)$, have been studied extensively, and have been measured in deep-inelastic scattering experiments [1]. The remaining distribution, $h_1(x)$, is relatively new, and has only recently begun to receive attention in the literature [2]. It has not been measured – since it is chiral-odd, it is inaccessible to inclusive deep-inelastic scattering experiments. Ralston and Soper [3] have shown, however, that $h_1(x)$ plays an important role in polarized Drell-Yan processes. More recently, Collins [4] showed that it emerges naturally in the factorization of a general hard process into soft and hard sub-processes. In the parton model language, $h_1(x)$ can be interpreted loosely as counting transversely polarized (valence) quarks in a transversely polarized nucleon.* The Q^2 evolution of $h_1(x)$ has been calculated by Kodaira *et al.* [5], and by Artru and Mekhfi [6].

Sum rules relating the moments of $h_1(x)$ to the nucleon matrix elements of local operators can be formulated in the standard way. The low moments are particularly interesting because they can be used to determine the gross features of the distribution. The lowest moment, for example, is referred to as the “tensor charge,” δq :

$$\delta q(Q^2) = \int_{-\infty}^{\infty} dx h_1(x, Q^2) = \int_0^1 dx \left(h_1(x, Q^2) - \bar{h}_1(x, Q^2) \right) . \quad (1)$$

Note that $h_1(x)$ carries flavor indices, which have been suppressed; there is an independent tensor charge, δq , for each quark flavor. The tensor charge derives its name from its relation to the following nucleon matrix element:

$$\langle PS | \bar{q} \sigma_{\mu\nu} i \gamma^5 q | PS \rangle \Big|_{Q^2} = \frac{2}{M} (S_\mu P_\nu - S_\nu P_\mu) \delta q(Q^2) \quad (2)$$

where $q = u, d, s$, *etc.* (The parameter Q^2 appearing in equations (1) and (2) is a renormalization scale label, necessary to render $h_1(x)$ and the tensor charges well-defined in perturbative QCD. The tensor charges do not mix with other operators under renormalization, and are therefore characterized by a single anomalous dimension, which has been calculated at one loop [5,6]: $\gamma = 2\alpha_s/3\pi$. Henceforth, we will suppress all dependence on Q^2 .) It is convenient to arrange the δq ’s into octet and singlet combinations, having definite flavor SU(3) transformation properties:

$$\begin{aligned} g_T^3 &= 2(\delta u - \delta d) \\ g_T^8 &= \frac{2}{\sqrt{3}}(\delta u + \delta d - 2\delta s) \\ g_T^0 &= 2\sqrt{\frac{2}{3}}(\delta u + \delta d + \delta s) . \end{aligned} \quad (3)$$

The reason for the peculiar normalization will become clear shortly. For future reference, we also define the axial charges, g_A^a :

$$\langle PS | \bar{\Psi} \lambda^a \gamma_\mu \gamma^5 \Psi | PS \rangle = S_\mu g_A^a . \quad (4)$$

* A more precise interpretation involves the transversely projected Pauli-Lubanski operator; see reference [2].

This is just the conventional definition of the axial charges. In the non-relativistic quark model, the tensor charges are equal to the corresponding axial charges. (We refer here to the octet and singlet combinations, g_T^a and g_A^a , *not* the individual quark components, δq , and their axial charge counterparts, Δq . According to the conventional definition of Δq , *i.e.*, $\langle PS|\bar{q}\gamma_\mu\gamma^5 q|PS\rangle = S_\mu\Delta q$, we find that $\Delta q = 2\delta q$, in the non-relativistic quark model. In this paper, we will work almost exclusively with g_T^a and g_A^a .) Jaffe and Ji [2] considered the tensor charges in the bag model, and found the tensor charges to be somewhat larger in absolute magnitude than the corresponding axial charges. It should be remembered, however, that this comparison involves an implicit, unknown scale, of the order of 1 GeV (*i.e.*, the “bag” scale at which the operators are renormalized).

The purpose of this note is to report the results of a Skyrme model analysis of the tensor charges. * In the large- N_c , SU(3)-symmetric limit, the model predicts that the octet isosinglet tensor charge, g_T^8 , is of order $1/N_c$ with respect to the (octet) isovector tensor charge, g_T^3 :

$$\frac{g_T^8}{g_T^3} = O\left(\frac{1}{N_c}\right) . \quad (5)$$

This can also be phrased in terms of a prediction for the F/D ratio:

$$\frac{F}{D} = \frac{1}{3} + O\left(\frac{1}{N_c}\right) . \quad (6)$$

These predictions coincide with the large- N_c , SU(3)-symmetric Skyrme model predictions for the octet *axial* charges, g_A^8 and g_A^3 . (The prediction $F/D = 1/3$ for the axial charges differs from the commonly quoted prediction of 5/9, which is based on an inconsistent treatment of the large- N_c limit.) The model also predicts that the singlet tensor charge, g_T^0 , is of order $1/N_c$ with respect to g_T^3 :

$$\frac{g_T^0}{g_T^3} = O\left(\frac{1}{N_c}\right) . \quad (7)$$

(In the case of the *axial* charges, the model predicts $g_A^0/g_A^3 = O(\frac{1}{N_c^2})$ [7].) These predictions are independent of the renormalization scale in QCD, since the tensor charges have a common anomalous dimension.

Before considering the Skyrme model, it will be useful to discuss the non-relativistic quark model (NQM). As mentioned above, the tensor charges are equal to the corresponding axial charges in the NQM. This is true for any value of N_c . The explicit calculation of the axial charges (and thus, the tensor charges) for arbitrary N_c has been carried out by Karl and Paton [8], using the fact that the axial charges measure the spin carried by the quarks. Transcribing their results, we have

$$\begin{aligned} g_T^3 &= \frac{1}{3}(N_c + 2) \\ g_T^0 &= \sqrt{2} g_T^8 = \sqrt{\frac{2}{3}} . \end{aligned} \quad (8)$$

* We work with the Skyrme model in order to be concrete. However, the conclusions are valid in the large- N_c limit of any chiral soliton model in which the soliton is quantized semi-classically, through the use of collective coordinates.

In particular, the NQM predicts that g_T^8/g_T^3 and g_T^0/g_T^3 are of order $1/N_c$. The known correspondence [9] between the large- N_c NQM and the large- N_c Skyrme model suggests that we should expect similar results in the Skyrme model.

The Skyrme model [10] describes the interactions of an octet of fundamental meson fields. In its simplest form, the effective action is given by

$$\Gamma = \int d^4x \left(\frac{F_\pi^2}{16} \text{Tr} (\partial_\mu U \partial^\mu U^\dagger) + \frac{1}{32e^2} \text{Tr} [(\partial_\mu U)U^\dagger, (\partial_\nu U)U^\dagger]^2 \right) + N_c \Gamma_{WZ} \quad , \quad (9)$$

where $U(x) = \exp(2i\lambda^a \phi^a(x)/F_\pi)$, and the ϕ_a 's are meson fields. F_π is the pion decay constant, and e is a phenomenological parameter. Γ_{WZ} is the so-called Wess-Zumino term, which incorporates the effects of anomalies. As is well known, the classical field equations associated with this lagrangian admit stable soliton solutions, the prototypical example of which is the hedgehog solution:

$$U_0(\vec{x}) = \begin{pmatrix} \exp(iF(r)\vec{\tau} \cdot \hat{x}) & 0 \\ 0 & 1 \end{pmatrix} \quad (10)$$

This is quantized by introducing collective coordinates to describe the orientation of the soliton in the group space $\text{SU}(3)_{\text{flavor}}$ (and thus, simultaneously, $\text{SU}(2)_{\text{spin}}$):

$$U(\vec{x}, t) = A(t)U_0(\vec{x})A^{-1}(t) \quad , \quad (11)$$

where $A(t)$ is a time-dependent $\text{SU}(3)$ matrix. The resulting quantum states have the quantum numbers of baryons. By making a suitable choice of the parameters F_π and e , the spectrum of low-lying baryons can be modeled approximately.

In order to apply the Skyrme model in estimating nucleon matrix elements, one must first identify the appropriate Skyrme model operators, constructed in terms of meson fields. The tensor charges are not associated with any symmetry of the QCD lagrangian, so their Skyrme model analogs cannot be generated by means of the standard Noether's theorem route. Other methods of constructing equivalent operators in the Skyrme model were considered, such as the method of Goldstone and Wilczek [11], but these apparently lead to ambiguous results. At leading order in the $1/N_c$ expansion, however, the analysis simplifies considerably, and we can learn about the flavor structure of the tensor charges simply by considering the transformation properties of the operators appearing in equation (2).

Consider the following matrix element:

$$\langle PS | \bar{\Psi} \lambda^a \sigma_{\mu\nu} i\gamma^5 \Psi | PS \rangle = \frac{1}{M} (S_\mu P_\nu - S_\nu P_\mu) g_T^a \quad . \quad (12)$$

For a nucleon at rest, the tensor operator appearing in this equation yields non-vanishing matrix elements only for $(\mu, \nu) = (i, 0)$, $i = 1, 2, 3$. Denote these components of the operator by O_i^a , and denote the equivalent Skyrme model operators by $\tilde{O}_i^a$. The $\tilde{O}_i^a$ transform as a three-vector under spatial rotations, and as an octet under $\text{SU}(3)$ transformations (or as an $\text{SU}(3)$ singlet, in the case of $a = 0$). In the large- N_c limit, the soliton rotates slowly, so as a first approximation, we ignore operators that involve time-derivatives, or in other words, factors of the canonical momenta. (The large- N_c approximation is standard in applications

of the Skyrme model; indeed, it is assumed implicitly in the quantization of the hedgehog solution.) Then in direct analogy with the quantum mechanics of a non-relativistic point particle, we can evaluate the matrix elements of the $\tilde{O}_i^a$ in terms of SU(3) Clebsch-Gordan coefficients, and a single (unknown) constant. Standard Skyrme model formulas [9] give the following results for the octet charges:

$$\langle N | \tilde{O}_m^a | N \rangle = k \sum_{n=1,2} \left(\begin{array}{c|c} \mathbf{R} & \mathbf{8} \\ N & a \end{array} \middle| \begin{array}{c} \mathbf{R}_n \\ N \end{array} \right) \left(\begin{array}{c|c} \mathbf{R} & \mathbf{8} \\ N & b \end{array} \middle| \begin{array}{c} \mathbf{R}_n \\ N \end{array} \right) \quad (13)$$

The label N represents a nucleon state, which we take to have definite polarization in the z -direction, for simplicity. The index m indicates the J_3 spin quantum number of the operator $\tilde{O}^a$. The label $\mathbf{R}$ denotes the SU(3) representation containing the nucleon – we will say more about this shortly. The direct product $\mathbf{R} \times \mathbf{8}$ contains two copies of the representation $\mathbf{R}$, which are denoted by $\mathbf{R}_n$ in equation (13). The label b represents a state with hypercharge zero, and isospin quantum numbers that are determined by the transformation properties of $\tilde{O}_m^a$ under spatial rotations: $I = J = 1$ and $I_3 = -J_3 = -m$.

The constant k is unknown, but the scale-dependence of the tensor charges in QCD means that precise knowledge of k would be of questionable value, since we do not know how to determine the renormalization scale at which k applies. Note that k may depend on N_c . Indeed, as we will see, correspondence with the NQM suggests that $k = O(N_c)$.

The corresponding formula for the singlet charge yields a vanishing result, since there is no state b having $I = 1$ in the singlet representation. It should be emphasized that equation (13) and its singlet counterpart are valid only when we neglect the contributions of operators involving time-derivatives. The corrections that result from such operators are discussed below. For now, we point out that these corrections do not necessarily vanish as N_c becomes large.

We now discuss the identification of the large- N_c analog of the nucleon. In a symmetric quark model, a baryon containing $N_c = 2n + 1$ quarks must belong to the $(1, n)$ representation of SU(3) if it is to have spin $\frac{1}{2}$. (The $(1, n)$ representation consists of the traceless, symmetric tensors having one upper index and n lower indices.) Furthermore, for $N_c = 2n + 1$, the $(1, n)$ representation is the only representation which can be projected out of the soliton solution and which contains states with zero strangeness and $I = J = \frac{1}{2}$ [9]. A consistent treatment of the large- N_c limit therefore seems to require that we choose $\mathbf{R} = (1, n)$ in equation (13). (For a more extensive discussion on this issue, see reference [12].) With this choice, equation (13) yields the following results:

$$\begin{aligned} g_T^3 &= 2k \frac{(N_c - 1)(N_c + 6)^2 + 9(N_c + 7)}{3(N_c + 7)(N_c + 3)^2} \xrightarrow{N_c \rightarrow \infty} \frac{2k}{3} \\ g_T^8 &= 2k \frac{2\sqrt{3}}{(N_c + 3)(N_c + 7)} \xrightarrow{N_c \rightarrow \infty} \frac{4\sqrt{3}k}{N_c^2} . \end{aligned} \quad (14)$$

Thus, the ratio g_T^8/g_T^3 vanishes in the large- N_c limit, although we will see that it is expected to vanish linearly in $1/N_c$, rather than quadratically. For comparison with the axial charges, it is useful to translate this result into a statement about the F/D ratio. According to the

standard definition of the reduced matrix elements, F and D , the isosinglet tensor charge is given by:

$$g_T^8 \sim \frac{1}{\sqrt{3}}(3F - D) \quad . \quad (15)$$

Thus, the large- N_c prediction is $F/D = 1/3$. As mentioned previously, this prediction is independent of the renormalization scale.

We now consider corrections to these results within the framework of the Skyrme model. Equation (13) is valid only if the operators that describe the tensor charges do not involve time-derivatives of the collective coordinates. Presumably, the quark operators appearing in equation (12) are matched onto an entire family of Skyrme model operators, and in general, this family will include operators with time-derivatives. We can argue on dimensional grounds, however, that the contributions of these operators are suppressed in the large- N_c limit. In the process of reducing meson operators to collective coordinate operators (by integrating over all space), a space-derivative brings in a mass parameter of the order of the inverse proton radius. In contrast, a time-derivative brings in a mass parameter of the order of the rotational kinetic energy of the soliton. Thus, we consider the ratio

$$\frac{L^2/2I}{1/R} \sim \frac{L^2}{MR} \quad (16)$$

where I is the moment of inertia, and L is the angular momentum of the spinning soliton. The nucleon mass grows as N_c , while the radius remains constant in the large- N_c limit [13]. Thus, we conclude that corrections arising from operators with time-derivatives should be suppressed by $1/N_c$.

In order to apply these ideas to the tensor charges, it is useful to consider the individual quark components, δu , δd , and δs . (This can be done unambiguously, since the octet and singlet tensor charges are characterized by the same anomalous dimension.) As suggested by the NQM, we will assume that δu and δd are of order N_c . Equation (14) then requires that δs be at most of order N_c . Recalling the considerations of the preceding paragraph, we find that operators with time-derivatives are expected to contribute to the δq 's at order unity. Equation (13) and its singlet counterpart then imply that g_T^3 is of order N_c , while g_T^0 and g_T^8 are of order unity (and thus, finite, in the large- N_c limit). These conclusions, of course, depend on the validity of our assumption that δu and δd are of order N_c . However, the conclusion that g_T^8/g_T^3 and g_T^0/g_T^3 are of order $1/N_c$ is readily seen to be independent of this assumption, and can be taken as a legitimate prediction of the Skyrme model. From $g_T^8/g_T^3 = O(\frac{1}{N_c})$, we also obtain $F/D = 1/3 + O(\frac{1}{N_c})$.

Now consider the axial charges. For a nucleon at rest, only the space-components of the axial-vector current operator appearing in equation (4) yield non-vanishing matrix elements. If we denote the space-components by A_i^a , then the A_i^a transform in exactly the same way as the O_i^a under $SU(3)_{\text{flavor}} \times SU(2)_{\text{spin}}$. Thus, the analysis outlined above also applies to the axial charges, and we draw the same conclusions as before: The ratio g_A^8/g_A^3 is of order $1/N_c$, and $F/D = 1/3 + O(\frac{1}{N_c})$; the ratio g_A^0/g_A^3 vanishes in the large- N_c limit, although, as mentioned previously, this ratio is actually expected to vanish like $(1/N_c)^2$, rather than $1/N_c$. (This last result does not contradict the arguments above, which are only meant to place crude limits on the large- N_c behavior.)

Note that the prediction $F/D = 1/3$ for the axial charges differs from the commonly quoted prediction of $5/9$ (see, for example, reference [14]). The latter prediction results from choosing $\mathbf{R}=8$ in equation (13). Although this value (*i.e.*, $5/9$) is in good agreement with the experimental result of 0.58 ± 0.05 [15], it is based on an inconsistent treatment of the large- N_c limit, so the agreement should perhaps be regarded as fortuitous. The proper prediction, $F/D = 1/3 + O(\frac{1}{N_c})$, clearly is not very precise for the realistic case of $N_c = 3$, although it is at least consistent with the experimental result. In the case of the tensor charges, where no experimental information is available, it would seem that the only reasonable prediction we can make is $F/D = 1/3 + O(\frac{1}{N_c})$.

The experimental result for the ratio g_A^0/g_A^3 is in fact small. Using the experimental values $g_A^0 = 0.098 \pm 0.076 \pm 0.113$ and $g_A^3 = 1.254 \pm 0.006$ [16], we obtain $g_A^0/g_A^3 = 0.078 \pm 0.109$. Unfortunately, this result does not offer a useful standard against which to compare the prediction $g_T^0/g_T^3 = O(\frac{1}{N_c})$, since g_A^0/g_A^3 is expected to be of order $O(\frac{1}{N_c^2})$.

In the above analysis, we stated that δs is at most of order unity with respect to δu and δd . In fact, since g_T^8 and g_T^0 are both of order $1/N_c$ with respect to g_T^3 , δs is actually of order $1/N_c$ with respect to δu and δd . (It also follows that $\delta u + \delta d$ is of order $1/N_c$ with respect to δu and δd , as in the NQM.) The same conclusion can be drawn in the case of the axial charges. If Δq is the axial charge counterpart of the tensor charge δq , then Δs is of order $1/N_c$ with respect to Δu and Δd . Since Δu and Δd are of order N_c [10] in the Skyrme model, Δs , the fraction of the nucleon spin carried by strange quarks, is of order unity, and not of order N_c , as previously reported [7]. This is consistent with the suppression of the $q\bar{q}$ sea that is expected on general grounds in large- N_c QCD [13]. (The previous prediction that $\Delta s = O(N_c)$ in the Skyrme model was based on an “ N_c -independent” F/D ratio of $5/9$. The argument used in that case breaks down for $F/D = 1/3$.)

Finally, we discuss the effects of SU(3) symmetry violation on the above results. The role of the strange quark in the nucleon is poorly understood in general. In the particular case of the tensor charges, it does not seem possible to say anything rigorous about the effects of a symmetry-breaking mass term for the strange quark. Within the framework of the Skyrme model, for example, we might approach this problem from the point of view of dimensional analysis. Unfortunately, dimensional analysis appears to be completely unreliable for SU(3) symmetry-breaking effects in the Skyrminion, since the dimensionful parameter that sets the scale for symmetry-breaking effects is of order 250 MeV (*i.e.*, less than the kaon mass, a typical measure of symmetry violation in the chiral lagrangian) [12]. We simply cannot rule out the possibility of large corrections stemming from SU(3) violation. For what it is worth, we point out that SU(3) appears to be a very good symmetry in the case of the axial charges, although again, the reasons for this are not well understood.

SUMMARY

We consider the nucleon tensor charges in the Skyrme model. We show that in the large- N_c , SU(3)-symmetric limit, the model predicts that g_T^8/g_T^3 is of order $1/N_c$. This is equivalent to the prediction $F/D = 1/3 + O(\frac{1}{N_c})$, which, unfortunately, is not very precise for the realistic case of $N_c = 3$. These predictions are identical to the Skyrme model predictions for the octet *axial* charges. The model also predicts that g_T^0/g_T^3 is of order $1/N_c$. All of these predictions are in agreement with the predictions of the NQM. Unlike the NQM, the Skyrme model does not readily indicate the large- N_c behavior of, say, g_T^3 , although it is not inconsistent to

assume that g_T^3 is of order N_c in the Skyrme model. We point out that the commonly quoted prediction for the axial charge F/D ratio (*i.e.*, 5/9) is based on an inconsistent treatment of the large- N_c limit. A consistent treatment yields the prediction quoted above, and shows that the strange quark contribution to the nucleon spin is of order unity in the large- N_c Skyrme model.

ACKNOWLEDGEMENTS

I thank Robert Jaffe for suggesting this work, and for many useful discussions. I would also like to thank Matthias Burkhardt and Eric Sather for useful discussions.

REFERENCES

- [1] For a review and references, see A. Manohar, UCSD/PTH 92-10 (1992).
- [2] For a review and references, see R. L. Jaffe and Xiangdong Ji, *Nucl. Phys.* **B375** (1992) 527.
- [3] J. Ralston and D. E. Soper, *Nucl. Phys.* **B152** (1979) 109.
- [4] J. Collins, Penn State Preprint, 1990.
- [5] J. Kodaira, S. Matsuda, K. Sasaki, and T. Uematsu, *Nucl. Phys.* **B159** (1979) 99.
- [6] X. Artru and M. Mekhfi, *Z. Physik* **C45** (1990) 669.
- [7] S. Brodsky, J. Ellis, and M. Karliner, *Phys. Lett.* **B206** (1988) 309.
- [8] G. Karl and J. E. Paton, *Phys. Rev.* **D30** (1984) 238.
- [9] A. Manohar, *Nucl. Phys.* **B248** (1984) 19.
- [10] E. Witten, *Nucl. Phys.* **B223** (1983) 422, 433; G. Adkins, C. Nappi, and E. Witten, *Nucl. Phys.* **B228** (1983) 433; E. Guadagnini, *Nucl. Phys.* **B236** (1984) 35.
- [11] J. Goldstone and F. Wilczek, *Phys. Rev. Lett.* **47** (1981) 986.
- [12] D. Kaplan and I. Klebanov, *Nucl. Phys.* **B335** (1990) 45.
- [13] E. Witten, *Nucl. Phys.* **B160** (1979) 57.
- [14] J. Bijnens, H. Sonoda, and M. Wise, *Phys. Lett.* **B140** (1984) 421.
- [15] R. L. Jaffe and A. Manohar, *Nucl. Phys.* **B337** (1990) 509.
- [16] J. Ashman *et al.*, *Nucl. Phys.* **B328** (1989) 1.